



Original article

Validation of the Brazilian version of the VERITAS-Pro scale to assess adherence to prophylactic regimens in hemophilia



Adriana Aparecida Ferreira^{a,b,*}, Isabel Cristina Gonçalves Leite^a,
Natalie August Duncan^c

^a Universidade Federal de Juiz de Fora UFJF, Juiz de Fora, MG, Brazil

^b Fundação Centro de Hematologia e Hemoterapia de Minas Gerais Hemominas, Juiz de Fora, MG, Brazil

^c Indiana Hemophilia and Thrombosis Center, Indianapolis, IN, USA

ARTICLE INFO

Article history:

Received 4 June 2017

Accepted 15 September 2017

Available online 26 November 2017

Keywords:

Hemophilia

Medication adherence

Questionnaire

Validation study

ABSTRACT

Introduction: Prophylaxis is the treatment of choice for patients with severe hemophilia. Low adherence may limit the effectiveness of the prophylactic regimen, thereby compromising outcomes.

Objective: The objective of this study was to validate the Brazilian version of the VERITAS-Pro prophylaxis adherence scale, originally an American questionnaire that can be answered by the individual responsible for prophylaxis as well as by an observer.

Methods: The scale has 24 questions divided into six subscales: Routine, Dosage, Plan, Remember, Skip and Communicate. Participants were recruited at a blood center in south-eastern Brazil for validation and reliability analyses. Validation measures included the results obtained using analog visual scales of adherence, interval between medication dispensed by the treatment center pharmacy and the percentage of recommended doses administered and infusions registered in the patients' logs.

Results: The study included 32 individuals responsible for prophylaxis and five observers. The internal consistency was very good for the VERITAS-Pro total score, excellent for the Remember, Skip and Communicate subscales, good for the Dosage subscale, and acceptable for the Routine and Plan subscales. Twelve participants answered the questionnaire on more than one occasion to evaluate reproducibility. The intraclass correlation coefficient was excellent. Regarding convergent validity, the VERITAS-Pro scores were moderately correlated with the global adherence scale and with infusion log records, but showed a weak correlation with pharmacy dispensation records.

* Corresponding author at: Fundação Centro De Hematologia E Hemoterapia De Minas Gerais (Hemominas), Barão de Cataguases S/N, Juiz de Fora, MG, Brazil.

E-mail address: adriana.hemato@gmail.com (A.A. Ferreira).

<https://doi.org/10.1016/j.bjhh.2017.09.002>

2531-1379/© 2017 Published by Elsevier Editora Ltda. on behalf of Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular.

Conclusion: The Brazilian version of VERITAS-Pro is a valid and reliable instrument, enabling the understanding of specific factors related to non-adherence and allowing targeted interventions for proper treatment.

© 2017 Published by Elsevier Editora Ltda. on behalf of Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular.

Introduction

Prophylaxis in hemophilia consists of regular infusions of clotting factor VIII or IX concentrate for a period longer than eight consecutive weeks in order to prevent bleeding.^{1,2} The benefits of prophylaxis include decreased frequency of bleeding episodes, decreased need for emergency room visits and hospitalizations, prevention of arthropathy, increased physical activity and school attendance, and improved academic performance.^{1,3–5} Prophylaxis reduces long-term morbidity, thus improving quality of life.^{6,7} The World Health Organization recommends prophylaxis as standard therapy for people with severe hemophilia.^{2,8}

Non-adherence to the prophylactic regimen may limit the effectiveness of treatment with less prevention of bleeding.⁹ The lack of standardized methods to assess adherence to hemophilia prophylaxis limits the understanding of factors that facilitate or hinder the therapeutic program. Lack of awareness of adherence as a determining factor in health outcomes can lead to a waste of human and economic resources as well as underutilization of available medications.^{10,11}

VERITAS-Pro is a questionnaire created in the United States, based on focus groups, to assess specific components of adherence as well as global adherence to the proposed prophylactic regimen.¹² The objective of this study was to describe the psychometric properties of the Brazilian version of the VERITAS-Pro, demonstrating its usefulness as an instrument for clinical practice and research.

Methods

Translation and adaptation

VERITAS-Pro, originally developed in English, was translated into Brazilian Portuguese according to international translation and adaptation guidelines.¹³ Two independent translations to Brazilian Portuguese were prepared by native Brazilians. There were no relevant differences between the translations. The two resulting versions were combined, corrected by experts, and translated back into English by an American translator without knowledge of the original document. The Brazilian version has 24 questions divided into six subscales: Time, Dose, Plan, Remember, Skip, and Communicate, just as in the original questionnaire. The questions were written in a way that made it possible for both the individual responsible for the patient's prophylaxis and an observer to respond. The answers are presented as five-point Likert scales ranging from 'never' to 'always'. An 'always' response reflects the best possible adherence for some items, and the worst for

others. For each item, a numerical classification was assigned to the Likert scale, giving one point to the response representing the best adherence and five points to the worst adherence. Possible scores of each subscale range from 4 to 20 points, and the total score of the instrument, from 24 to 120 points, where 120 represents the worst adherence.

Participants

Participants were recruited at the *Hemocentro Regional de Juiz de Fora (HRJF)* in southeastern Brazil. The individuals considered eligible were those responsible for the prophylaxis of patients with hemophilia with a severe phenotype, A or B, in a home infusion regimen for at least six months. Patients in on-demand treatment, those not qualified to receive home infusions, and those with inhibitors were excluded.

Participant recruitment and data collection

The study was approved beforehand by the Research Ethics Committee of the *Fundação Centro de Hematologia e Hemoterapia de Minas Gerais – HEMOMINAS*. The participants were asked to respond to the questionnaire during their usual visits to the blood center. All signed a free and informed consent form, allowing access to their infusion logs, and received a guarantee of confidentiality of any individual information. Parents gave their consent to include data from patients under 18 years of age, and adolescent patients (12–17 years old) gave their agreement to participate. Sociodemographic and health data were collected from the patients, as well as sociodemographic data from those responsible for prophylaxis, in the case of under 18-year-old and incapacitated patients. Participants answered a questionnaire and a visual analog scale on global adherence to prophylaxis. For the test–retest evaluation, participants were asked to respond the questionnaire a second time on their next visit to the blood center. Data collection occurred between October 2015 and November 2016.

Other measures of adherence to prophylaxis

Analog scale of global adherence to prophylaxis

After completing the VERITAS-Pro, participants were reminded of the details of the prescribed prophylaxis. An analog scale with values ranging from zero (never follow the prescription) to 10 (always follow the prescription) was then presented and the participants were asked to rate their adherence in the previous three months.

Records of medications dispensed by the blood center pharmacy

The number of doses dispensed was divided by the number of doses required to comply with the prescribed prophylaxis in the interval between dispensations. The analysis included dispensations that provided sufficient factor for the previous twelve weeks of treatment. If the individuals responsible returned later than the expected time for taking the required doses, it became evident that prophylactic infusions were skipped. However, if they returned at shorter than expected intervals, as it was not possible to determine by this method the reason for consumption greater than that prescribed, the highest possible adherence was considered 100%.

Percentage of recommended doses administered

The percentage of doses administered on the correct days in relation to the doses prescribed in the three months prior to the application of the VERITAS-Pro was calculated from the infusion logs filled out by those responsible for the patient's prophylaxis. When clearly documented, doses to treat bleeding events were excluded. However, extra prophylactic doses infused prior to medical and dental procedures, or unusual physical activity, were included. Thus, the prophylaxis measured by this method could be greater than 100%, but for analytical statistics reasons, 100% was considered the maximum possible adherence value.

Statistical analysis

Reliability and validity

Internal consistency was assessed using Cronbach's alpha. Since the subscales of the instrument contain only four items each, the following considerations were used: 0.8 characterized excellent internal consistency; 0.7, very good; 0.6, good; and >0.5, minimally acceptable.¹² The intraclass correlation coefficient (ICC) was used to evaluate the VERITAS-Pro scores in the test-retest reliability analysis, which was considered excellent when >0.75.¹³ The Spearman correlation test was used to compare VERITAS-Pro results with the other adherence measures.¹⁴ Statistical analysis was carried out using the Statistical Package for the Social Sciences (SPSS) v.20 software.

Results

Participants

At the beginning of the study, in October 2015, there were 59 patients eligible for the prophylaxis regimen at the HRJF. Twenty-three patients (39%) had opted to remain on episodic treatment and one tested positive for an inhibitor (1.7%). Of the 36 patients who adhered to the prophylactic treatment, one was receiving the infusions at the blood center and one presented severe clinical complications, requiring frequent changes in the infusion scheme and, for these reasons, they were excluded from the study. Thus, those responsible for the prophylaxis of 34 patients and five observers were recruited for the research, but two patients declined participation. The 32 patients, whose prophylaxis adherence was evaluated, were all males, 27 diagnosed with hemophilia A

Table 1 – Sociodemographic characteristics of patients with hemophilia in a home prophylaxis regimen between October 2015 and November 2016.

Sociodemographic characteristic	
Age (years) – mean (range)	20.6 (3–60)
Self-declared color – n (%)	
White	20 (62.5)
Black	6 (18.8)
Mulatto	4 (12.5)
Indigenous	2 (6.3)
Marital status – n (%)	
Single	28 (87.5)
Married	2 (6.3)
Divorced	2 (6.3)
Schooling of over 18-year olds – n (%)	
Higher education completed or in progress	9 (56.4)
High school diploma	1 (6.3)
Complete elementary education	1 (6.3)
Incomplete elementary education	4 (25)
No education at school	1 (6.3)
Occupation of those over 18 – n (%)	
Student	7 (43.8)
Retired due to disability	6 (37.5)
Formal employee	2 (12.5)
Unemployed	1 (6.3)
Occupation of those under 18 – n (%)	
Student	15 (93.8)
No occupation	1 (6.2)

(21 with factor VIII dosage <1%) and five with hemophilia B (four with factor IX dosage <1%). The patients' ages ranged from three to 60 years (mean of 20.6 years, with a standard deviation of 14.1). All the boys under the age of six were regularly attending preschool. The patient was responsible for the infusions in only 11 cases (34.3%). Tables 1 and 2 show the sociodemographic and clinical characteristics of the patients, respectively.

VERITAS-Pro scores

The total mean of the VERITAS-Pro was 35.51 (range: 24–70) when the responses of those responsible for prophylaxis and the observers were analyzed together. There was no statistically significant difference when the scores were evaluated separately (35.47 and 35.8, respectively; p -value = 0.914). There was also no difference when the patients themselves were the primary infusers (38.91) or others took on this responsibility (33.81; p -value = 0.17). The averages of the subscales ranged from 4.59 (Dose) to 7.3 (Time) (Table 3).

VERITAS-Pro reliability

Results showed a very good internal consistency for the VERITAS-Pro total score; excellent for the Remember, Skip and Communicate subscales; good for the Dose subscale; and minimally acceptable for the Time and Plan subscales (Table 4).

Table 2 – Clinical characteristics and therapeutic aspects of hemophilia patients between October 2015 and November 2016.

Clinical characteristic	
Type of hemophilia – n (%)	
A	27 (84.4)
B	5 (15.6)
Severity – n (%)	
<1% Factor VIII or IX	25 (78.1)
1–2%b Factor VIII or IX	7 (21.9)
Family history of hemophilia – n (%)	
Yes	18 (56.3)
No	14 (43.8)
Presence of target joints before prophylaxis started – n (%)	
Yes	28 (87.5)
No	4 (12.5)
Clinically evident arthropathy – n (%)	
Yes	18 (56.3)
No	14 (43.8)
Physiotherapy in the past three months – n (%)	
Yes	5 (15.6)
No	27 (84.4)
Engage in regular physical activity – n (%)	
Yes	20 (62.5)
No	12 (37.5)
Time in prophylaxis – median (range)	21.6 months (6.5–40.6)
Spontaneous hemarthrosis in the past three months – n (%)	
None	20 (62.5)
1 or 2	9 (28.2)
3 or more	3 (9.4)
Primary infuser – n (%)	
Patient	11 (34.4)
Mother/father	8 (25)
Sister/brother	1 (3.1)
Other relative	1 (3.1)
Health professional	11 (34.4)

VERITAS-Pro reproducibility

Twelve participants (37.5%) agreed to answer the VERITAS-Pro questionnaire on more than one occasion for test–retest reliability analysis. The mean interval between the two responses was 55.5 days (standard deviation of 11.9 days; range: 24–145

Table 4 – Internal consistency analysis of the Brazilian version of the VERITAS-Pro.

Subscales	Cronbach's alpha	Cronbach's alpha if the item is deleted	Item-total correlation
Total	0.737		
Time	0.589	0.736	0.653
Dose	0.656	0.746	0.673
Plan	0.514	0.766	0.402
Remember	0.843	0.709	0.838
Skip	0.864	0.706	0.824
Communicate	0.826	0.731	0.635

Table 5 – Evaluation of reproducibility of the Brazilian version of the VERITAS-Pro.

Subscale	Test	Retest	Intraclass correlation coefficient	p-Value
Total	33.58	32.75	0.933	<0.01
Time	6.83	6.42	0.895	<0.01
Dose	4.75	4.75	0.986	<0.01
Plan	5.42	5.17	0.536	0.11
Remember	5.58	5.75	0.929	<0.01
Skip	5.08	5.67	0.234	0.33
Communicate	5.92	5.00	0.835	<0.01

days). The ICC for the test–retest was excellent for the total score (0.933; p -value <0.01), and for all subscales except for the Plan and Skip domains (Table 5).

Adherence assessed by infusion logs

The infusion logs of 32 patients were evaluated regarding the 12 weeks that preceded the application of the VERITAS-Pro. In all cases it was not possible to identify prophylactic infusions done outside the schedule of episodic infusions for the treatment of hemorrhagic events. Those responsible reported an average infusion rate of 72.17% of the prophylactic doses prescribed in the period (range: 25–105.5%; standard deviation: 25.8%). Half of those responsible for prophylaxis reported infusion of 80% or more of the prescribed doses,

Table 3 – VERITAS-Pro total and subscale scores.

	All participants			Responsible			Observers		
	Mean	Range	SD	Mean	Range	SD	Mean	Range	SD
Total	35.51	24–70	10.39	35.47	24–70	11.04	35.80	28–41	5.07
Time	7.30	4–14	2.47	7.19	4–14	2.50	8	6–11	2.34
Dose	4.59	4–12	1.67	4.59	4–12	1.73	4.60	4–7	1.34
Plan	5.51	4–13	2.06	5.47	4–13	2.17	5.80	4–7	1.30
Remember	5.57	4–14	2.55	5.56	4–14	2.67	5.60	4–8	1.81
Skip	6.08	4–16	2.77	6.06	4–16	2.87	6.20	4–9	2.28
Communicate	6.46	4–20	3.77	6.59	4–20	3.94	5.60	4–10	2.60

SD: standard deviation.

which characterizes good adherence, when this is evaluated dichotomously.^{7,10,15,16}

Adherence measured by visual analog scale

The global adherence reported by 35 participants (31 responsible for infusions and four observers) through the analog scale ranged from 6 to 10. Mean global adherence was 9.2 with a standard deviation of 1.05 and a median of 10.

Adherence measured by the blood center pharmacy's dispensing of factor concentrate

The blood center pharmacy usually dispenses doses sufficient for four weeks of prophylaxis on each visit of those responsible for infusions. Thus, the individuals responsible are expected to return with the used bottles at the end of this period and replenish their household supply. The dispensing of doses sufficient for prophylactic infusions in the 12 weeks prior to the survey was analyzed, and the actual time until the return of all used bottles was computed. Adherence was calculated by dividing the number of doses dispensed by the number of doses required to meet the prescription until the return, revealing a mean adherence of 89.02% (standard deviation of 18.3, with a minimum of 45.9% and a maximum of 130%).

Concurrent validity of the VERITAS-Pro

Since better adherence is characterized by lower scores on the VERITAS-Pro, higher scores on the visual analog scale, and by higher adherence percentages based on pharmacy dispensing and on infusion logs, negative correlations indicate a stronger correspondence of VERITAS-Pro with other adherence measures. Therefore, VERITAS-Pro scores were moderately correlated with the global adherence scale ($r = -0.529$; p -value = 0.002) and with the records in infusion logs ($r = -0.516$; p -value = 0.003), but showed a weak correlation with the pharmacy dispensing records ($r = -0.32$; p -value = 0.074).

Discussion

Prophylaxis is considered the standard therapy for patients with severe hemophilia¹⁷ as it is capable of preventing arthropathy when begun early,¹⁸ and of reducing the number of bleeding episodes, and improving the quality of life of individuals who already have irreversible joint damage.¹⁹ However, adherence to treatment is essential to achieve these results.^{20,21}

In Brazil, the prophylactic treatment regimens for hemophilia were incorporated by the Brazilian National Health System as of November 2011.²² The lack of a reliable instrument to assess adherence to prophylaxis has been one of the obstacles to research in this field since then.²³ To meet this need, the psychometric properties of the Brazilian version of the VERITAS-Pro were evaluated, showing good reliability and validity when applied to both those responsible for prophylaxis and to observers of the treatment carried out.

The total score showed a very good internal consistency, although lower than the original questionnaire ($\alpha = 0.737$ versus $\alpha = 0.92$). In addition, the scores had a moderate correlation with the visual scale of adherence and with the percentage of administered infusions recorded in the patients' logs. That is, the Brazilian version of the VERITAS-Pro is a measure with temporal stability and that relates with other measures of adherence to prophylaxis. Almost all the subscales showed good internal consistency, except for the Time ($\alpha = 0.589$) and Plan subscales ($\alpha = 0.514$). Particular attention should be paid to these subscales and items considering these subscales in future analysis because their modification could improve the quality of the scale. Unfortunately, a factorial analysis was not possible due to the size of the sample.

The total score and all the subscales showed excellent test-retest reliability, except for the Skip subscale (ICC = 0.234), whose means increased significantly in the participant's second assessment. This increase, which translates as worsening adherence in the Skip domain, was unexpected since individuals tend to improve their behavior, albeit transiently, after being evaluated for the first time.²⁴

As in the original study, the Dose subscale exhibited the lowest mean while the Time subscale presented the highest, indicating that participants reported better adherence in administering the correct dose and worse in administering it on prescribed days and times.¹²

In addition, as in the American work, the dispensing of factor concentrates by the pharmacy was a less useful validation measure than the infusion logs and the visual scale of adherence.¹² This is explained by the fact that the dispensing of medications includes variables that are not directly related to patient adherence, such as the healthcare model of the local health system. Cuesta-Barriuso et al. argue that free treatment can weaken the patient's commitment in following through with the prescriptions, but not in the acquisition of the medication.²⁵ While the infusion logs represent an excellent source of data for validation and are commonly used to monitor infusions at home, they only provide quantitative data.^{3,7,15,26}

The superiority of the VERITAS-Pro over the global measures is due to its ability to recognize the multiple facets of the adherence construct, represented by the subscales, and to perceive different types of behavior. While the Time, Dose and Skip subscales identify whether and when infusions were administered, the Plan and Communicate domains reveal the individual's baseline behavior. This more comprehensive view provided by the VERITAS-Pro assists in directing interventions to improve adherence to prophylaxis.¹²

Another advantage of the Brazilian version of the VERITAS-Pro was that it is easy to understand and quick to complete, making it suitable for routine use in the clinical practice.

It is believed that this is the first work in Brazil on adherence to prophylaxis in hemophilia. Although small, the sample was sufficient for the validation analyses of the Brazilian version of the VERITAS-Pro. Unfortunately, subgroup analyses were limited by the low number of participants. Although differences in adherence were not described according to the type or severity of hemophilia, or regarding the presence of infections such as hepatitis C or human immunodeficiency virus,²⁵ Miebach

and Kalnins reported that adherence varies according to age group and is lower among young adults.²⁷ Studies with larger populations are needed for these evaluations in the Brazilian context.

Finally, it should be noted that although all the children with severe hemophilia were in prophylaxis during this study, the situation of the adults was quite different. The results of this study showed that patients in the Brazilian sample had higher adherence to treatment than those enrolled in the study conducted by Duncan et al.,¹² but a significant percentage chose to continue with episodic treatment, expressing the lowest possible level of adherence to prophylaxis, yet these patients were not considered by the VERITAS-Pro.

Conclusion

The Brazilian version of the VERITAS-Pro is a quantitative and detailed instrument for assessing adherence to hemophilia prophylaxis. Evidence from this study supports its validity and reliability for use in the clinical and research contexts. This validated scale can aid in the understanding of behaviors and in the identification of modifiable determinants of non-adherence associated with unsatisfactory orthopedic outcomes. The findings derived from the VERITAS-Pro can facilitate the design of targeted interventions in these situations, but future studies are needed to confirm the relationship between adherence, patient characteristics, and clinical outcomes.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

The authors thank Maria Cristina Belletti Rodrigues for her assistance with one of the translations of the original questionnaire.

REFERENCES

- Shapiro AD, Donfield SM, Lynn HS, Cool VA, Stehbins JA, Hunsberger SL, et al. Defining the impact of hemophilia: the academic achievement in children with hemophilia study. *Pediatrics*. 2001;108(6):e105.
- National Hemophilia Foundation. Medical and Scientific Advisory Council (MASAC) Recommendations Concerning Prophylaxis. MASAC Document 179. New York, NY: National Hemophilia Foundation; 2007. p. 10001.
- Manco-Johnson MJ, Abshire TC, Shapiro AD, Riske B, Hacker MR, Kilcoyne R, et al. Prophylaxis versus episodic treatment to prevent joint disease in boys with severe hemophilia. *N Engl J Med*. 2007;357(6):535-44.
- Löfqvist T, Nilsson IM, Berntorp E, Pettersson H. Haemophilia prophylaxis in young patients – a long-term follow-up. *J Intern Med*. 1997;241(5):395-400.
- Fischer K, van der Bom JG, Molho P, Negrier C, Mauser-Bunschoten EP, Roosendaal G, et al. Prophylactic versus on-demand treatment strategies for severe haemophilia: a comparison of costs and long-term outcome. *Haemophilia*. 2002;8(6):745-52.
- Panicker J, Warriar I, Thomas R, Lusher J. The overall effectiveness of prophylaxis in severe haemophilia. *Haemophilia*. 2003;9(3):272-8.
- Thornburg C, Pipe S. Adherence to prophylactic infusions of factor VIII or factor IX for haemophilia. *Haemophilia*. 2006;12(2):198-9.
- World Health Organization. Delivery of Treatment for Haemophilia: Report of a Joint WHO/WFH/ISTH Meeting London, UK 11-13 February 2002. London, UK: WHO; 2002.
- Hacker M, Geraghty S, Manco-Johnson M. Barriers to compliance with prophylaxis therapy in haemophilia. *Haemophilia*. 2001;7(4):392-6.
- Thornburg C. Physicians' perceptions of adherence to prophylactic clotting factor infusions. *Haemophilia*. 2008;14(1):25-9.
- Shapiro A. Why is primary prophylaxis underutilized in the United States? *Haemophilia*. 2003;9(6):670-2.
- Duncan N, Kronenberger W, Roberson C, Shapiro A. VERITAS-Pro: a new measure of adherence to prophylactic regimens in haemophilia. *Haemophilia*. 2010;16(2):247-55.
- Acquadro C, Conway K, Hareendran A, Aaronson N. European Regulatory Issues and Quality of Life Assessment (ERIQA) Group. Literature review of methods to translate health-related quality of life questionnaires for use in multinational clinical trials. *Value Health*. 2008;11(3):509-21.
- Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet*. 1986;1(8476):307-10.
- du Treil S, Rice J, Leissing C. Quantifying adherence to treatment and its relationship to quality of life in a well-characterized haemophilia population. *Haemophilia*. 2007;13(5):493-501.
- Lam WY, Fresco P. Medication adherence measures: an overview. *BioMed Res Int*. 2015;2015:217047.
- Srivastava A, Brewer AK, Mauser-Bunschoten EP, Key NS, Kitchen S, Llinas A, et al. Treatment Guidelines Working Group on Behalf of The World Federation of Hemophilia. Guidelines for the management of hemophilia. *Haemophilia*. 2013;19(1):e1-47.
- Oldenburg J. Optimal treatment strategies for hemophilia: achievements and limitations of current prophylactic regimens. *Blood*. 2015;125(13):2038-44.
- Manco-Johnson MJ, Kempton CL, Reding MT, Lissitchkov T, Goranov S, Gercheva L, et al. Randomized, controlled, parallel-group trial of routine prophylaxis vs. on-demand treatment with sucrose-formulated recombinant factor VIII in adults with severe hemophilia A (SPINART). *J Thromb Haemost*. 2013;11(6):1119-27.
- Brown MT, Bussel JK. Medication adherence: WHO cares? *Mayo Clin Proc*. 2011;86(4):304-14.
- Krishnan S, Vietri J, Furlan R, Duncan N. Adherence to prophylaxis is associated with better outcomes in moderate and severe haemophilia: results of a patient survey. *Haemophilia*. 2015;21(1):64-70.
- Brasil. Ministério da Saúde. SUS garante tratamento preventivo para hemofílicos. Brasília: Ministério da Saúde; 2011. Available from: <http://portalsaude.saude.gov.br/index.php/cidadao/principal/agencia-saude/noticias-antecedentes-agencia-saude/1229-sus-garante-tratamento-br-preventivo-para-hemofílicos> [cited 16.12.11].
- Lock J, Raat H, Duncan N, Shapiro A, Beijlevelt M, Peters M, et al. Adherence to treatment in a Western European paediatric population with haemophilia: reliability and

- validity of the VERITAS-Pro scale. *Haemophilia*. 2014;20(5):616-23.
24. Miller L, Lovler R, McIntire S. *Foundations of psychological testing: a practical approach*. 4th ed. Los Angeles: SAGE Publication, Inc.; 2013. p. 156-89 [Chapter 6: What is test reliability?].
25. Cuesta-Barriuso R, Torres-Ortuño A, Galindo-Piñana P, Nieto-Munuera J, Duncan N, López-Pina JA. Validation of the VERITAS-Pro treatment adherence scale in a Spanish sample population with hemophilia. *Patient Prefer Adher*. 2017;11:653-60.
26. Tencer T, Roberson C, Duncan N, Johnson K, Shapiro A. A haemophilia treatment center-administered disease management programme in patients with bleeding disorders. *Haemophilia*. 2007;13(5):480-8.
27. Miesbach W, Kalnins W. Adherence to prophylactic treatment in patients with haemophilia in Germany. *Haemophilia*. 2016;22(5):e367-74.